RETICULITERMES FLAVIPES (ISOPTERA: RHINOTERMITIDAE) and DROSOPHILA MELANOGASTER (DIPTERA: DROSOPHILIDAE) SHARE DEVELOPMENTAL GENE SEQUENCE

Tracie M. Jenkins¹, Tea Meulia², Susan C. Jones³, Brian T. Forschler⁴, and Saskia A. Hogenhout⁵

¹⁾Department of Entomology, University of Georgia, Griffin, Georgia
²⁾Molecular and Cellular Imaging Center, Ohio State University, OARDC, Wooster, Ohio
³⁾Department of Entomology, Ohio State University, Columbus, Ohio
⁴⁾Department of Entomology, University of Georgia, Athens, Georgia
⁵⁾Department of Entomology, Ohio State University, OARDC, Wooster, Ohio

Key Words Subterranean termites eyelid gene gene sequence

After constructing a *Reticulitermes flavipes* (Kollar) worker complementary DNA (cDNA) library and screening clones, we found a gene fragment homologous to the eyelid (*eld*) gene of *Drosophila melanogaster*. The *eld* gene is required for embryonic segmentation, development of the notum and wing margin, and photoreceptor differentiation in *D. melanogaster*. We designed oligonucleotide primers based on termite and *Drosophila* sequences, then used these primers to amplify and sequence through the gene fragment. We report the first characterization of the *eld* gene sequence in a subterranean termite.